



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Area deprivation, screen time and consumption of food and drink high in fat salt and sugar (HFSS) in young people: Results from a cross sectional study in the UK

Citation for published version:

Thomas, F, Thomas, C, Hooper, L, Rosenberg, G, Vohra, J & Bauld, L 2019, 'Area deprivation, screen time and consumption of food and drink high in fat salt and sugar (HFSS) in young people: Results from a cross sectional study in the UK', BMJ Open. <https://doi.org/10.1136/bmjopen-2018-027333>

Digital Object Identifier (DOI):

[10.1136/bmjopen-2018-027333](https://doi.org/10.1136/bmjopen-2018-027333)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMJ Open

Publisher Rights Statement:

This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



BMJ Open Area deprivation, screen time and consumption of food and drink high in fat salt and sugar (HFSS) in young people: results from a cross-sectional study in the UK

Fiona Thomas,¹ Christopher Thomas,¹ Lucie Hooper,¹ Gillian Rosenberg,¹ Jyotsna Vohra,¹ Linda Bauld²

To cite: Thomas F, Thomas C, Hooper L, *et al.* Area deprivation, screen time and consumption of food and drink high in fat salt and sugar (HFSS) in young people: results from a cross-sectional study in the UK. *BMJ Open* 2019;**9**:e027333. doi:10.1136/bmjopen-2018-027333

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-027333>).

Received 17 October 2018
Revised 6 June 2019
Accepted 7 June 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

¹Cancer Policy Research Centre, Cancer Research UK, London, UK

²Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK

Correspondence to

Miss Fiona Thomas;
fiona.thomas@hotmail.com

ABSTRACT

Objectives To investigate associations between deprivation in young people and consumption of foods high in fat, salt and sugar (HFSS), screen time exposure and health knowledge.

Design An online cross-sectional survey with people aged 11–19 years in the UK, where participants reported consumption behaviours across 13 HFSS and two non-HFSS groups; screen time for commercial television and streaming services; and knowledge of health conditions and their links to obesity.

Setting UK

Participants A total of 3348 young people aged 11–19 years across the UK.

Main outcome measures The study assessed the consumption behaviours, commercial screen time exposure and the health knowledge of 3348 people aged 11–19 years. Multivariate binary regression analysis, controlling for age and gender, was performed.

Results Deprivation level was associated with increases in consumption of six of the HFSS products including energy drinks (OR: 2.943, $p<0.001$) and sugary drinks (OR: 1.938, $p<0.001$) and a reduction in consumption in the two non-HFSS products included in the study, fruit (OR: 0.668, $p=0.004$) and vegetables (OR: 0.306, $p<0.001$). Deprivation was associated with high weekly screen time of both television (OR: 2.477, $p<0.001$) and streaming (OR: 1.679, $p=0.001$). Health knowledge was also associated with deprivation. There was lower awareness of the association of obesity and cancer (OR: 0.697, $p=0.003$), type 2 diabetes (OR: 0.64, $p=0.004$) and heart disease (OR: 0.519, $p<0.001$) in the most deprived.

Conclusions Young people from the more deprived areas of the UK were more likely to consume a range of HFSS products, report increased exposure to HFSS advertising and have a poorer awareness of health conditions associated with overweight and obesity. The findings suggest that population-level measures addressing childhood obesity should account for consumption patterns among different groups of children and young people and the factors that may influence these.

Strengths and limitations of this study

- The study identified an association between socioeconomic deprivation and risk factors that may influence the prevalence of childhood obesity in the UK through a nationally representative sample of young people aged 11–19 years across England and each of the devolved nations.
- Data collected for this study were from a single cross-sectional survey, so it is not possible to determine causation between the variables.
- The study could not directly assess exposure to marketing of foods HFSS, although previous research that had explored the relationship between commercial screen time and exposure and found the two to be related provided us with greater confidence that screen time may be a reliable proxy for marketing exposure under current UK marketing regulations.

BACKGROUND

In the UK, around 30% of children are overweight or obese,¹ the highest rate of childhood obesity in Europe. This overall figure masks considerable disparities by socioeconomic status. Overweight and obesity prevalence for children in the 10% most deprived areas in England, for example, are more than double that of those who live in the least deprived 10%.² Longer term, an obese child is around five times more likely to become an obese adult,³ and there is substantial evidence that obesity in adulthood directly contributes to the development of conditions such as diabetes, coronary heart disease, stroke and 13 different types of cancer.^{4–8}

Previous studies have provided diverse explanations for the rise in levels of obesity, ranging from genetics, increased calorie intake, an increase in sedentary behaviour or

a combination of factors.^{9–11} The calorie intake increase is thought to be the most significant influence accounting for this rise, caused by a range of environmental factors,^{9 12 13} including the role of the marketing and promotion of foods that are high in fat, salt and sugar (HFSS), often referred to as ‘junk foods’. Marketing of these foods is extensive and delivered through a variety of platforms including television, streaming, price promotions and print media. Studies have identified a substantial expenditure by manufacturers and retailers on the marketing of junk food to children and young adults^{14 15} and identified that industry recognises the potential marketing has for influencing consumption choices.¹⁶ The link between marketing and weight outcomes, as well as increased consumption of HFSS products, has been highlighted by a number of previous studies.^{17–23} Assessing exposure to HFSS marketing via self-reported recall of viewing advertisements does have limitations. Thus, some studies have used commercial screen time as a proxy for TV and online marketing exposure,^{17 21} whereby greater screen time indicates increased exposure to HFSS advertising. Prior content analysis of UK television, where young people make up a large proportion of the audience, highlighted the increased likelihood of HFSS marketing exposure,²⁴ supporting this proxy measure.

Increasingly, there is a need to identify how marketing and promotion affect children in different social groups or those living in more or less deprived communities. Studies have previously identified an association between socioeconomic status and obesity.^{25 26} This association is likely to be caused by a wide variety of factors including the pricing and availability of particular products in a locality,²⁷ understanding and awareness of dietary factors and weight and social norms.^{28 29} However, further research is required to understand the inter-relationship of these factors and also other drivers of consumption including exposure to HFSS advertising. This type of research is important to identify interventions or policy actions that can contribute to addressing overweight and obesity. Halting or reversing current obesity trends is a current priority for public health policies in the UK,^{30 31} and globally.^{32 33}

Hooper *et al*³⁴ identified that there is a low level of public awareness of the link between overweight and obesity, resulting in preventable health conditions, including cancer. Only 25% of the UK adult population are aware of this link, and this lack of awareness is more prevalent in less affluent groups. Other studies have also found an association between greater health awareness and increased support for policy change, particularly for alcohol policy.^{35 36} Greater health knowledge may therefore affect how young people view the acceptability of HFSS marketing and also consumption choices.

To date, there is limited research on the association between deprivation, HFSS marketing and obesity. Given these gaps, this study aims to investigate whether such a relationship exists and how it might be influenced by particular mediators such as frequency and duration

of exposure to marketing and knowledge and understanding of health risks.

METHODS

Study design

An online cross-sectional survey was conducted between April and June 2017. The survey was developed following cognitive testing with a small sample of young people (n=100) to ensure age and cultural comprehensibility of the questions, some of which were based on well-validated questions used in other surveys.^{21 37–40} This survey was compiled using the prevalidated questions and the advice of senior researchers from the National Cancer Institute, USA and Public Health England and the comprehensive guidance of the Institute for Social Marketing and the University of Stirling who have experience of running the Youth Tobacco Policy Survey. The final survey covered six main themes: exercise levels, food and drink consumption, screen time, recalled marketing exposure, perceptions of marketing and demographic factors.

A total of 3348 young people, aged 11–19 years, were recruited by market research company, YouGov, using their in-house panel. YouGov already had data on the children in households of adult in-house panel members. Children over the age of 16 years were directly approached and asked if they wished to participate. For those aged under 16 years, their parents were contacted and asked if their child could participate in the survey. Data collected were weighted by age, gender, ethnicity, region and social grade to be representative of the UK population.

Measures

Deprivation

Level of deprivation was assessed using an area-based measure rather than individual measure of socio-economic status. The Index of Multiple Deprivation (IMD) was coded into five equal quintiles for analysis, ranging from 1, the lowest 20% of deprivation, to 5, the highest 20% of respondents. IMD is a measure of the relative deprivation of an area, combining information from seven domains: income deprivation, employment deprivation, education, skills and training deprivation, health deprivation and disability, crime, barriers to housing and services and living environment deprivation.⁴¹

Consumption behaviours

The survey measured consumption of a range of food and drink products. Participants were asked to report their consumption behaviours from the question ‘How often do you usually eat or drink...?’ followed by a series of food categories, HFSS including biscuits and cakes, chips, confectionary, crisps, desserts, energy drinks, flavoured yoghourts, milk-based drinks, ready meals, sugary drinks, sweetened cereal and takeaways, as well as with healthy items such as fruit and vegetables. These food groups were chosen using previous research on unhealthy products and with reference to the categories

included by Public Health England in their sugar reduction programme⁴² and the nutrient profiling model to ascertain nutritional composition of foods. Responses were graded on a Likert scale from more than once a day to never and then converted to binary variables across two coding groups. The groups identified as 'higher' consumption depending on the total calorific content in each food.^{17 21 42} The first coding group included sugar-sweetened drinks, flavoured yoghourts, confectionary, cakes/biscuits, fruit, vegetables, diet drinks, crisps and desserts where two or more portions a week was considered high consumption.⁴² The second coding group included takeaways, ready meals, energy drinks, fried potato products, milk-based drinks and sugar-sweetened cereals where one or more portions a week was considered higher consumption.⁴² This coding was deduced from a range of approaches including the Public Health England nutrient model, pilot testing and a review of the average calories in each product, recognising the differences between portion sizes of the food categories. The coding was only calculated for participants who gave an answer, and those who selected 'not sure' were excluded from the final analysis.

Screen time

Commercial screen time was a variable created in the data set based on responses related to frequency and duration of exposure to TV and streaming (on demand) services.^{21 43} Participants listed the hours spent watching both commercial and non-commercial TV and streaming services. This excluded screen time from computers being used for homework. Non-commercial screen time (which contains no paid for marketing in the UK context) was shown to not be significant in previous analysis of the data^{17–19} and was therefore removed. Weekend and week-day viewing was then weighted and turned into a weekly measure and categorised; low (<3 hours per week), medium (3–21 hours per week) and high (21 hours or more per week).²¹

Health knowledge

Health knowledge was assessed using the question 'Which, if any, of the following health conditions do you think can result from being overweight? Please tick all that apply.' Options included answers that were both correct and incorrect to identify the extent of health knowledge. The eight chosen conditions were cancer, stroke, heart disease, diabetes type 1, diabetes type 2, migraines, chicken pox and influenza. From this, the results were coded as a binary variable: 0—unaware; 1—aware of the links between certain conditions and being overweight.

Age and gender

Control variables were selected on theoretical importance from a rapid review of the literature^{21 40 43–46} and included gender (coded 0—male, 1—female) and age (11–19 years).

Patient and public involvement

Prior to the development of the survey, qualitative research was carried out by colleagues at the University of Stirling and National Centre for Social Research. This work consulted young people through focus groups, on the design and content of the questionnaire. Results of these focus group discussions were published in a Cancer Research UK report 'It's Just There To Trick You'.⁴⁷ This Patient and Public Involvement (PPI) development work involved discussion of relevant research questions related to food and beverage consumption, relevant policy issues (ie, exposure to food marketing, pricing, availability), use of broadcast media examples and the appropriateness of questions relating to sociodemographic characteristics. Questions included in the resulting survey were then trialled with young people using cognitive interviewing techniques as described above. Survey reports are publicly available on Cancer Policy Research Centre's website.

ANALYSIS

Data were analysed using IBM SPSS Version 23. Multiple multivariate binary regression models were run on the unweighted data to test for associations between deprivation levels and three key behaviours of young people—consumption behaviours, screen time use and health knowledge. The consumption model used the dependent binary variables of food and drink consumption behaviours. Models were run separately for each dependent variable, producing 15 models in total. The screen time model used the dependent variable of categorised reported screen time hours. The health knowledge model used the dependent binary variable of awareness of a health condition and its link to overweight and obesity. This included eight different health conditions, some with identified associations, and some without.

Within each of these models, the IMD variable was used as an independent variable, with the least deprived quintile as the reference group. Age and gender were included in the models as control variables, as potentially confounding variables.

RESULTS

Sample characteristics

Almost half (49%) of the survey respondents were female and 51% male. The mean age of participants was aged 14.9 years (SD=2.55). The majority (82%) were from white British backgrounds with 18% from other ethnic groups. The majority of respondents lived in England (82%); 5% of respondents lived in Wales, 8% in Scotland and 3% in Northern Ireland (table 1).

Screen time

Of respondents, there were 31.9% in low, 57.1% in medium and 11.0% in high screen time categories for television viewing. For streaming services, there was

Table 1 Sample demographics of the UK representative respondents

Male	
11–12	11.0%
13–15	16.0%
16–17	12.0%
18–19	12.0%
Female	
11–12	10.0%
13–15	16.0%
16–17	11.0%
18–19	12.0%
Ethnicity	
White	82.0%
BME	18.0%
IMD	
1,2	20.0%
3,4	20.0%
5,6	20.0%
7,8	20.0%
9,10	20.0%
Region	
North East	4.0%
North West	11.1%
Yorkshire & Humber	8.5%
East Midlands	7.3%
West Midlands	9.3%
East	9.3%
London	12.7%
South East	14.0%
South West	8.2%
Wales	4.7%
Scotland	7.8%
Northern Ireland	3.1%

BME, Black and Minority Ethnicity; IMD, Index of Multiple Deprivation.

30.3% in low, 50.7% in medium and 19.0% in high screen time categories.

Deprivation and consumption behaviours

The results of the binary logistic regressions showed an association between deprivation and higher consumption behaviours, for a range of HFSS food products (table 2). The most deprived young people were significantly more likely to consume energy drinks (OR=2.943, $p<0.001$), followed by sugary drinks (OR=1.938, $p<0.001$).

In contrast, analysis identified consumption of fruit and vegetables was inversely associated with more deprived groups. Fruit (OR=0.668, $p=0.004$) and vegetables

(OR=0.306, $p<0.001$) were more likely to be consumed in lower frequency by the most deprived respondents, when compared with the most affluent respondents, as per the use of IMD. Therefore, these young people had a reduced likelihood of consuming the healthier options in higher quantities.

Deprivation and screen time

Regression analysis found an association between deprivation in young people and high weekly screen time of both television and streaming (table 3). The model compared 'high' category screen time (21 hours or more a week) to 'medium and low' screen time (<21 hours a week) and found those from the most deprived quintile were significantly more likely to be in the high screen time category than the more affluent respondents, for both television (OR=2.477, $p<0.001$) and streaming (OR=1.679, $p=0.001$).

Deprivation and health knowledge

The analysis identified an association between deprivation and poorer health knowledge (table 4). Respondents were asked whether eight health conditions (from a pre-existing list) could occur as a result of being overweight or obese. There was significantly poorer awareness of the association between cancer (OR=0.697, $p=0.003$), type 2 diabetes (OR=0.64, $p=0.004$) and heart disease (OR=0.519, $p<0.001$) and obesity for those from the more deprived quintiles. There was also significantly higher association between incorrectly linking type 1 diabetes (OR=1.536, $p<0.001$) and obesity in the most deprived quintile, compared with the most affluent quintile.

DISCUSSION

Results from this survey identify a clear association between socioeconomic deprivation and risk factors that may influence the prevalence of childhood obesity in the UK. Involving a nationally representative sample of young people aged 11–19 years across England and each of the devolved nations, it sought to explore whether young people living in more deprived areas reported knowledge and behaviours that may contribute to obesity. The study found that these young people consumed more foods and beverages high in salt sugar and fat and were conversely less likely to report consumption of fruit and vegetables. In addition, young people living in more deprived communities spent more time watching commercial broadcast media where they could be exposed to HFSS advertising. Young people living in less affluent areas also had lower levels of awareness of the preventable health conditions, including cancer, which can arise as a result of obesity.

These results support findings from previous studies on factors influencing childhood obesity but also provide new evidence on the clustering of these factors among less affluent groups. It is well established that

Table 2 Consumption behaviours and deprivation

	Consumption behaviours				Logistic regression/significance		
	Descriptive findings				Most deprived quintile		
	Most deprived 20%		Least deprived 20%				
	High consumption (%)	Low consumption (%)	High consumption (%)	Low consumption (%)	OR	CI (95%)	value
Biscuits and cakes	58.4	41.6	63.6	36.4	0.841	0.638 to 1.038	0.097
Chips	72.1	27.9	67.5	32.5	1.259	0.974 to 1.627	0.079
Confectionary	65.3	34.7	66.3	33.7	0.962	0.751 to 1.232	0.759
Crisps	62.8	37.2	58.5	41.5	1.232	0.965 to 1.572	0.093
Desserts	47.1	52.9	55.2	44.8	0.732	0.576 to 0.930	0.011
Diet drinks	35.7	64.3	31.2	68.8	1.233	0.959 to 1.585	0.102
Energy drinks	15.6	84.4	7.0	93.0	2.493	1.676 to 3.706	0.000
Flavoured yoghourts	28.0	72.0	27.3	72.7	1.061	0.812 to 1.386	0.067
Fruit	71.3	28.7	78.8	21.2	0.668	0.507 to 0.879	0.004
Milk-based drinks	31.3	68.7	22.6	77.4	1.613	1.229 to 2.118	0.001
Ready meals	64.3	35.7	56.3	43.7	1.416	1.111 to 1.712	0.005
Sugary drinks	41.6	58.4	27.2	72.8	1.938	1.506 to 2.494	0.000
Sweetened cereals	49.6	50.4	44.8	55.2	1.253	0.986 to 1.593	0.066
Takeaways	39.1	60.9	25.1	74.9	1.914	1.482 to 2.472	0.000
Vegetables	78.9	21.1	92.4	7.6	0.306	0.211 to 0.442	0.000

The bolded values indicate significance from the model.

there is a clear gradient in overweight and obesity by socioeconomic status in both adults and young people, with individuals from less affluent communities more likely to carry excess weight compared with their more affluent neighbours.^{2 25 26 48–50} More limited research has explored how eating patterns vary by deprivation in young people. This study adds to existing evidence, suggesting that greater HFSS consumption and lower levels of fruit and vegetable consumption are more common in less affluent young people.^{18 51}

This survey also asked young people about the time they spend watching television and on-demand screening services and calculated ‘screen time’ using

an approach employed in previous studies.^{17 21} While higher levels of screen time are associated with sedentary behaviour, which may contribute to obesity, they may also suggest greater exposure to broadcast media marketing including of HFSS foods. Previous research has found that children who spent more time watching commercial TV and on demand programmes in the UK are exposed to more HFSS food marketing than those with lower levels of screen time.^{24 52} Viewing more HFSS advertisements on TV and streaming has been associated with higher HFSS consumption, with the difference between a high consumer and a low consumer being at least 520 junk food products a year.¹⁷

Table 3 Screen time behaviour and deprivation

	Screen time behaviours						Logistic regression/significance		
	Frequency						Most deprived quintile		
	Most deprived (20%)			Least deprived (20%)					
	Low viewing (%)	Medium viewing (%)	High viewing (%)	Low viewing (%)	Medium viewing (%)	High viewing (%)	OR	CI	p value
Television screen time	25.7	55.9	18.4	31.7	60.0	8.3	2.477	1.697–3.614	0.000
Streaming screen time	28.2	46.8	25.0	33.2	50.3	16.5	1.679	1.234–2.283	0.001

The bolded values indicate significance from the model.

Table 4 Health knowledge and deprivation

	Overall awareness	Frequency		Logistic regression/significance	
		Most deprived awareness %	Least deprived awareness %	OR	p value
Cancer link	42.0	36.1	44.4	0.697	[0.003]
Heart disease link	87.0	83.5	90.3	0.519	[0.000]
Stroke link	60.0	62.1	59.2	0.862	[0.228]
Diabetes type 1 link	39.0	46.1	36.3	1.536	[0.000]
Diabetes type 2 link	82.0	78.7	84.7	0.64	[0.004]
Influenza link	4.0	4.1	2.7	1.544	[0.196]
Chicken pox link	1.0	0.4	0.7	0.558	[0.502]
Migraine link	14.0	11.0	11.8	0.907	[0.604]

The bolded values indicate significance from the model.

We also found that young people living in more deprived areas had lower levels of awareness of the links between overweight and obesity and relevant health conditions. Awareness is relevant because evidence relating to other preventable risk factors (such as smoking and alcohol) suggest that health knowledge is relevant as a preliminary step toward changing behaviour, but also, importantly, understanding and support for policies and interventions that may address key factors that drive consumption including restrictions on marketing and pricing of unhealthy products.^{35 36 53 54}

Taken together, our findings suggest that inequalities in rates of obesity in young people in the UK may be linked to knowledge and behaviours driven by key aspects of an obesogenic environment. Action to address childhood obesity needs to take into account differential consumption patterns among less affluent young people and the factors that may influence these consumption patterns. The introduction of policies and interventions that aim to address these factors, including better information on the health consequences of obesity, reducing exposure to HFSS marketing and other wider population level measures (such as policies to address the price and content of products) should consider and assess their impact on less affluent groups.

Strengths and limitations of the study

This study has a number of limitations. Data are from a single cross-sectional survey, so it is not possible to determine causation. The measure of deprivation used was an area-based measure (IMD), which has limitations, as individual or household deprivation may vary within areas. Young people may have limited knowledge and awareness of different health conditions, for example, the distinction between type 1 and type 2 diabetes, and therefore, responses to the question relating to health conditions may in part reflect this lack of understanding. Responses to each of the key topics of interest including food consumption patterns, use of streaming services and TV viewing,

and awareness of health conditions linked to obesity were based on self-report and thus subject to misreporting or recall bias. The study could not directly assess exposure to HFSS marketing, although our previous research has explored the relationship between commercial screen time and exposure and found the two to be related,^{17 21} which provides us with greater confidence that screen time may be a reliable proxy for marketing exposure under current UK marketing regulations. The response rate for this survey is estimated at 26% for people aged 16–19 years and 47% for people aged 11–15 years with the parental consent process. This is in line with what is usually obtained for surveys with young people by YouGov and is a limitation when sampling from this population demographic. Finally, overweight and obesity in young people are driven by a wide range of factors beyond those assessed in this study.

Conclusions and future research

Taken together our findings suggest that inequalities in rates of obesity in young people in the UK may be linked to knowledge and behaviours driven by key aspects of an obesogenic environment. Action to address childhood obesity needs to take into account differential consumption patterns among less affluent young people and the factors that may influence these consumption patterns. The introduction of policies and interventions that aim to address these factors, including better information on the health consequences of obesity, reducing exposure to HFSS marketing and other wider population level measures (such as policies to address the price and content of products) should consider and assess their impact on less affluent groups.

Future research should explore in more detail a larger number of factors, including, for example, the affordability and availability of HFSS foods, social norms and the influence of social networks in more deprived communities and how these influence knowledge and behaviour among more deprived young people. In addition, studies should assess the impact of changes to the

policy and regulatory environment proposed in the UK and other countries to reduce childhood obesity and how these changes may affect young people living in communities where obesity rates are highest.

Acknowledgements The authors are grateful to the Cancer Policy Research Centre, Cancer Research UK for funding the study. The views expressed are those of the researchers and not necessarily those of the funder.

Contributors Fiona Thomas carried out the majority of the analysis for this publication along with Chris Thomas and Lucie Hooper who also developed the questions for the survey. Gillian Rosenberg contributed to the preparation of the manuscript and analysis plan. Both Jyotsna Vohra and Linda Bauld contributed to the study design and concept and the preparation of the manuscript.

Funding This study was funded by Cancer Research UK (<http://dx.doi.org/10.13039/501100000289>).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethical approval was obtained in January 2017 from the General University Ethics Panel at the University of Stirling. This ethical approval covered both cognitive testing of the questionnaires and the online surveys. YouGov's staff included lead for ethical and quality assurance to ensure adherence to best practice throughout testing and data collection.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement There is no additional unpublished data available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

1. National Child Measurement Programme (NCMP). *Trends in child BMI*: Public Health England, 2017.
2. Patterns and trends in child obesity. 2018 <https://www.gov.uk/guidance/phe-data-and-analysis-tools#obesity-diet-and-physical-activity2018>.
3. Simmonds M, Llewellyn A, Owen CG, et al. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* 2016;17:95–107.
4. Parkin DM, Boyd L. 8. Cancers attributable to overweight and obesity in the UK in 2010. *Br J Cancer* 2011;105(Suppl 2):S34–7.
5. Ades PA, Savage PD. Obesity in coronary heart disease: An unaddressed behavioral risk factor. *Prev Med* 2017;104:117–9.
6. Brown KF, Rungby H, Dunlop C, et al. The fraction of cancer attributable to modifiable risk factors in England, Wales, Scotland, Northern Ireland, and the United Kingdom in 2015. *Br J Cancer* 2018;118:1130–41.
7. Public Health England. *Adult obesity and type 2 diabetes*, 2014.
8. World Health Organisation. Global Health Risks-Mortality and burden of disease attributable to selected major risks. *The Lancet* 2015.
9. Bleich S, Cutler D, Murray C, et al. Why is the developed world obese? *Annu Rev Public Health* 2008;29:273–95.
10. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet* 2002;360:473–82.
11. Reilly JJ. Obesity in childhood and adolescence: evidence based clinical and public health perspectives. *Postgrad Med J* 2006;82:429–37.
12. Afshin A, Peñalvo JL, Del Gobbo L, et al. The prospective impact of food pricing on improving dietary consumption: A systematic review and meta-analysis. *PLoS One* 2017;12:e0172277.
13. Vandevijvere S, Chow CC, Hall KD, et al. Increased food energy supply as a major driver of the obesity epidemic: a global analysis. *Bull World Health Organ* 2015;93:446–56.
14. Obesity Health Alliance. *Health costs of obesity soaring as junk food companies pour millions into advertising*, 2017.
15. Powell LM, Harris JL, Fox T. Food marketing expenditures aimed at youth: putting the numbers in context. *Am J Prev Med* 2013;45:453–61.
16. Story M, French S. Food advertising and marketing directed at children and adolescents in the US. *Int J Behav Nutr Phys Act* 2004;1:3.
17. Thomas C, Hooper L, Petty R, et al. *10 years on*: Cancer Research UK, 2018.
18. Thomas C, Hooper L, Petty R, et al. *Under Pressure: New evidence on young people's broadcast marketing exposure in the UK*: Cancer Research UK, 2018.
19. Thomas F, Hooper L, Petty R, et al. *A prime time for action: new evidence on the link between television and on-demand marketing and obesity*: Cancer Research UK, 2018.
20. Boyland EJ, Nolan S, Kelly B, et al. Advertising as a cue to consume: a systematic review and meta-analysis of the effects of acute exposure to unhealthy food and nonalcoholic beverage advertising on intake in children and adults. *Am J Clin Nutr* 2016;103:519–33.
21. Scully M, Wakefield M, Niven P, et al. Association between food marketing exposure and adolescents' food choices and eating behaviors. *Appetite* 2012;58:1–5.
22. Halford JC, Boyland EJ, Hughes G, et al. Beyond-brand effect of television (TV) food advertisements/commercials on caloric intake and food choice of 5–7-year-old children. *Appetite* 2007;49:263–7.
23. National Child Measurement Programme. *Changes in children's body mass index between 2006/07 and 2015/16*: Public Health England, 2017.
24. Boyland EJ, Harrold JA, Kirkham TC, et al. The extent of food advertising to children on UK television in 2008. *Int J Pediatr Obes* 2011;6:455–61.
25. Wang Y. Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. *Int J Epidemiol* 2001;30:1129–36.
26. Public Health England. Mills J, ed. *Severe obesity in ten to eleven year olds reaches record high*: Public Health England, 2018.
27. Howard Wilsher S, Harrison F, Yamoah F, et al. The relationship between unhealthy food sales, socio-economic deprivation and childhood weight status: results of a cross-sectional study in England. *Int J Behav Nutr Phys Act* 2016;13:21.
28. Sanderson SC, Waller J, Jarvis MJ, et al. Awareness of lifestyle risk factors for cancer and heart disease among adults in the UK. *Patient Educ Couns* 2009;74:221–7.
29. Puhl R, Suh Y. Health consequences of weight stigma: implications for obesity prevention and treatment. *Curr Obes Rep* 2015;4:182–90.
30. Scottish Government. *A Healthier Future - Scotland's diet and healthy weight delivery plan*. Edinburgh: Scottish Government, 2018.
31. Department of Health and Social Care. Chapter 2: HM Government. *Global public health directorate: obesity FaN. Childhood Obesity: a plan for action*, 2018.
32. Swinburn BA. Obesity prevention: the role of policies, laws and regulations. *Aust New Zealand Health Policy* 2008;5:12.
33. WHO. World health organisation: obesity and overweight. 2017 <http://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight> (Accessed 24 Jul 2018).
34. Hooper L, Anderson AS, Birch J, et al. Public awareness and healthcare professional advice for obesity as a risk factor for cancer in the UK: a cross-sectional survey. *J Public Health* 2018;40:797–805.
35. Buykx P, Li J, Lovatt M, et al. *An examination of public attitudes towards alcohol policy*: University of Sheffield and Cancer Research UK, 2016.
36. Bates S, Holmes J, Gavens L, et al. Awareness of alcohol as a risk factor for cancer is associated with public support for alcohol policies. *BMC Public Health* 2018;18:688.
37. Public Health England. National diet and nutrition survey. 2016 <https://www.gov.uk/government/collections/national-diet-and-nutrition-survey> (Accessed 03 Jan 2018).
38. Marketing IoS. *Food choice survey*: University of Stirling, 2007.
39. Alcohol (Minimum Unit) Pricing Act. 2012 http://www.legislation.gov.uk/asp/2012/4/pdfs/asp_20120004_en.pdf.
40. Morley BC, Scully ML, Niven PH, et al. What factors are associated with excess body weight in Australian secondary school students? *Med J Aust* 2012;196:189–92.
41. Department for Communities and Local Government. *The English Index of Multiple Deprivation (IMD) - Guidance*, 2015.
42. Public Health England. *Sugar reduction: achieving the 20%*, 2017.
43. Dixon HG, Scully ML, Wakefield MA, et al. The effects of television advertisements for junk food versus nutritious food on children's food attitudes and preferences. *Soc Sci Med* 2007;65:1311–23.
44. Denny S, Lewycka S, Utter J, et al. The association between socioeconomic deprivation and secondary school students' health: findings from a latent class analysis of a national adolescent health survey. *Int J Equity Health* 2016;15:109.

45. Nestle M. Food marketing and childhood obesity--a matter of policy. *N Engl J Med* 2006;354:2527-9.
46. Cooke LJ, Wardle J. Age and gender differences in children's food preferences. *Br J Nutr* 2005;93:741-6.
47. MacGregor A, Biquelet A, H L, et al. "It's just there to trick you": A qualitative study of 11-19 year olds perceptions of food and drink marketing.": ScotCen Social Research; NatCen Social Research; Institute for Social Marketing, University of Stirling; and Policy Research Centre for Cancer Prevention, Cancer Research UK, 2016.
48. Stamatakis E, Wardle J, Cole TJ. Childhood obesity and overweight prevalence trends in England: evidence for growing socioeconomic disparities. *Int J Obes* 2010;34:41-7.
49. McLaren L. Socioeconomic status and obesity. *Epidemiol Rev* 2007;29:29-48.
50. Bann D, Johnson W, Li L, et al. Socioeconomic inequalities in body mass index across adulthood: coordinated analyses of individual participant data from three british birth cohort studies initiated in 1946, 1958 and 1970. *PLoS Med* 2017;14:e1002214.
51. Crawford B, Byun R, Mitchell E, et al. Socioeconomic differences in the cost, availability and quality of healthy food in Sydney. *Aust N Z J Public Health* 2017;41:567-71.
52. Campbell D. Children seeing up to 12 adverts for junk food an hour on TV, study finds. 2017 <https://www.theguardian.com/society/2017/nov/28/children-seeing-up-to-12-adverts-for-junk-food-an-hour-on-tv-study-finds> (Accessed 23 Jul 2018).
53. Bryant T. Role of knowledge in public health and health promotion policy change. *Health Promot Int* 2002;17:89-98.
54. Raphael D, Bryant T. The state's role in promoting population health: public health concerns in Canada, USA, UK, and Sweden. *Health Policy* 2006;78:39-55.